



SPECIAL EXAMINATION **MARCH 2024**

COURSE TITLE

STATISTICS FOR SOCIAL SCIENCE

COURSE CODE

RMAT2233

DATE / DAY

11 JULY 2024 / THURSDAY

TIME/DURATION

09:00 AM - 12:00 PM / 03 Hour(s) 00 Minute(s)

INSTRUCTIONS TO CAL

1. Please read the instruction under each section carefully

Candidates are reminded not to bring into examination hall/room any form of written materials or electronic 2. gadget except for stationery that is permitted by the Invigilator.

Students who are caught breaching the Examination Rules and Regulation will be charged with an academic 3. dishonesty and if found guilty of the offence, the maximum penalty is expulsion from the University.

(This Question Paper consists of 8 Printed Pages including front page)

Copying, modifying, or reprinting, is not permitted.

There are SEVEN (7) questions in this section. Answer ALL questions in the answer booklet provided. [100 MARKS]

1.There are 15 Malays, 10 Indians and x Chinese students in a Statistics class. If the value of x for each of the following:	Find the
a) if the probability of choosing Malay is .	(5 marks)
b) if the probability of choosing an Indian student is .	(5 marks)
 A stall sells onions. The weights of onions are normally distributed with a mea 85 grams and standard deviation 5 grams. Five onions are chosen at random probability 	
a) exactly three of them weigh more than 82 grams.	(5 marks)
b) at least one onion is more than 82 grams.	(5 marks)
3.The table below shows the years of experience of 120 employees of Uni Raza	k.
Years of experience Number of employees	
Tours of employees	-
Of Modis	
Wing, or, WAD	2
eprinting is not permitted	
Calculate	
a) Mean.	(5 marks)
b) Median.	(5 marks)
c) Mode.	(5 marks)
I. For the followings:	
a) Find the skew of this data set:	
Calculate Pearson's Coefficient of Skewness for above sample	(5 marks)
b) Find the standard deviation of the sample data below:	5 marks)

5. A certain type of ball is known to have a bounce height which is normally is



5. A certain type of ball is known to have a bounce height which is normally is normally distributed with a standard deviation of 2 cm. A random sample of 60 tennis ball is tested and the mean bounce height of the sample is 140 cm. Find

a) a symmetrical 95% confidence interval for mean bounce height.

(10 marks)

b) a symmetrical 99% confidence interval for mean bounce height.

(10 marks)

The weights of durian in Dusun Gombak is normally distributed with a mean of 5 kg and and a standard deviation of 0.5 kg. Given there are 100 durians in a basket find the number of durians that

a) weigh more than 6kg.

(10 marks)

b) weigh more than 4 kg.

(10 marks)

7. The masses of guavas in a farm are normally distributed with a mean μ and a standard deviation, σ . The mass of percentages of guava that less than 400 g is 15.87% and more than 500 g is 6.68%. Find the value of μ and σ . (15 marks)

Copying, matter END OF QUESTION PAPER ***

ABDUL RAZAL

RAZAL

RAZAL

Copying, modifying, or reprinting, is not permitted.

List of Formulas

- 1. Sample Mean: $\bar{X} = \frac{\sum x}{n}$
- 2. Population Mean: $\mu = \frac{\sum X}{N}$

3. Grouped Data

Mean:
$$\vec{X} = \frac{\sum fx}{\sum f}$$

$$Median = L_m + \left[\frac{\frac{n}{2} - F}{f_m}\right] c$$

$$Mode = L + \left[\frac{f_0 - f_1}{(f_0 - f_1) + (f_0 - f_2)} \right] \times c$$

4. Population (Ungrouped Data)

$$Mean: \mu = \frac{\sum x}{N}$$

Variance:
$$\sigma^2 = \frac{\sum x^2}{N} - (\bar{X})^2 @ \frac{1}{N} \sum (X - \mu)^2$$

Standard deviation:
$$\sigma = \sqrt{\frac{\sum x^2}{N} - (\bar{X})^2}$$
 @ $\sqrt{\frac{1}{N} \sum (X - \mu)^2}$

5.Sample (Ungrouped Data)

Variance:
$$s^2 = \frac{1}{n-1} \left[\sum x^2 - \frac{(\sum x)^2}{n} \right]$$

Standard deviation:
$$s = \sqrt{\frac{1}{n-1} \left[\sum x^2 - \frac{(\sum x)^2}{n} \right]}$$

6. Population (Grouped Data)

$$Mean: \mu = \frac{\sum fx}{\sum f(N)}$$

Variance:
$$\sigma^2 = \frac{\sum fx^2}{\sum f(N)} - (\bar{X})^2$$

Standard deviation:
$$\sigma = \sqrt{\frac{\sum fx^2}{\sum f(N)} - (\bar{X})^2}$$
.

7. Sample (Grouped Data)

Variance:
$$s^2 = \frac{1}{n-1} \left[\sum_{j=1}^{n} fx^2 - \frac{(\sum_{j=1}^{n} fx)^2}{n} \right]$$

Standard deviation:
$$s = \sqrt{\frac{1}{n-1} \left[\sum fx^2 - \frac{(\sum fx)^2}{n} \right]}$$

8. Pearson's coefficient of skewness

$$Skewness = \frac{Mean - Mode}{Standard\ deviation}\ @\ \frac{3(\ Mean - Median)}{Standard\ deviation}$$

Copying, modifying, or reprinting, is not permitted.

9.Binomial

$$P(X=r) = C_r p^r q^{n-r}$$

Mean = np

Variance = npq

Standard deviation = \sqrt{npq}

10. Poisson Distribution

$$P(X=r)=e^{-\mu}\frac{\mu^r}{r!}$$

11.Normal Distribution

$$z = \frac{X - \mu}{\sigma}$$

12.

$$E(\bar{X}) = \mu$$

$$Var\left(\bar{X}\right) = \frac{\sigma^2}{n}$$

13.Confidence Interval for Population Mean

(with known variance & sample size > 30)

$$P(\bar{X}-E \leq \mu \leq \bar{X}+E),$$

$$E = \pm z_{\frac{\alpha}{2}} \frac{\sigma}{\sqrt{n}}, E = marginal of error$$

Confidence interval will

$$(\bar{X}-E,\bar{X}+E)$$

ABDUL RAZ, 14.Confidence Interval for Population Mean (with unknown variance & sample size < 30)

$$(\bar{X} - E \leq \bar{X} \leq \bar{X} + E)$$

$$E = \pm t_{\frac{\alpha}{2}} \frac{\hat{\sigma}}{\sqrt{n}}$$

Confidence interval will

$$(\bar{X}-t_{\frac{\alpha}{2}}\frac{\partial}{\sqrt{n}},\bar{X}+t_{\frac{\alpha}{2}}\frac{\partial}{\sqrt{n}})$$



15.

Significance test

Population mean (Normal) with known variance

Test Statistics
$$z = \frac{\overline{x} - \mu}{\frac{\sigma}{\sqrt{n}}}$$

Population mean (Normal) with unknown variance

Test Statistics
$$z = \frac{x - \mu}{\frac{S}{\sqrt{n}}}$$

16.Anova

Test Statistics:
$$\frac{{s_1}^2}{{s_2}^2}$$

 $F = \frac{estimated\ population\ variance\ between\ the\ sample}{estimated\ population\ variance\ within\ the\ sample}$

17.

$$r = \frac{n(\sum XY) - (\sum X)(\sum Y)}{\sqrt{[n\sum X^2 - (\sum X)^2][n\sum Y^2 - (\sum Y)^2]}}$$

Test statistics:
$$t = \frac{r\sqrt{n-2}}{\sqrt{1-r^2}}$$

$$b = \frac{n\sum XY - (\sum X)(\sum Y)}{n(\sum X^2) - (\sum X)^2} \quad a = \frac{\sum Y}{n} - b\frac{\sum X}{n}$$

$$s_{y.x} = \sqrt{\frac{\sum (Y - Y')^2}{n - 2}}$$

$$Test statistics, t = \frac{b}{SE(b)}$$

18.

Confidence Interval of an Estimate

$$Y' \pm t_{\frac{\alpha}{2}} s_{y,x} \sqrt{\frac{1}{n} + \frac{(X - \overline{X})^2}{\sum X^2 - \left[\frac{(\sum X)^2}{n}\right]}}$$


Prediction Interval of an Estimate

$$Y' \pm t_{\frac{\alpha}{2}} s_{y.x} \sqrt{1 + \frac{1}{n} + \frac{(X - \bar{X})^2}{\sum X^2 - \left[\frac{(\sum X)^2}{n}\right]}}$$



THE UPPER TAIL PROBABILITY Q(z) FOR THE NORMAL DISTRIBUTION N(0,1) KEBARANGKALIAN HUJUNG ATAS Q(z) BAGI TABURAN NORMAL N(0, 1)

2	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
1	u l		-	J	•	,	ď	f		,		Minus / Tolak							
0.0	0.5000	0.4960	0.4920	0.4880	0.4840	0.4801	0.4761	0.4721	0.4681	0.4641	4	8	12	16	20	24	28	32	36
0.1	0.4602	0.4562	0.4522	0.4483	0.4443	0.4404	0.4364	0.4325	0.4286	0.4247	4	8	12	16	20	24	28	32	36
0.2	0.4207	0.4168	0.4129	0.4090	0.4052	0.4013	0.3974	0.3936	0.3897	0.3859	4	8	12	15	19	23	27	31	35
0.3	0.3821	0.3783	0.3745	0.3707	0.3669	0.3632	0.3594	0.3557	0.3520	0.3483	4	7	11	15	19	22	26	30	34
0.4	0.3446	0.3409	0.3372	0.3336	0.3300	0.3264	0.3228	0.3192	0.3156	0.3121	4	7	11	15	18	22	25	29	32
0.5	0.3085	0.3050	0.3015	0,2981	0.2946	0.2912	0.2877	0.2843	0.2810	0.2776	3	7	10	14	17	20	24	27	31
0.6	0.2743	0.2709	0.2676	0.2643	0.2611	0.2578	0.2546	0.2514	0.2483	0.2451	3	7	10	13	16	19	23	26	29
0.7	0.2420	0.2389	0.2358	0.2327	0.2296	0.2266	0.2236	0.2206	0.2177	0.2148	3	6	9	12	15	18	21	24	27
0.8	0.2119	0.2090	0.2061	0.2033	0.2005	0.1977	0.1949	0.1922	0.1894	0.1867	3	5	8	11	14	16	19	22	25
0.9	0.1841	0.1814	0.1788	0.1762	0.1736	0.1711	0.1685	0.1660	0.1635	0.1611	3	5	8	10	13	15	18	20	23
1.0	0.1587	0.1562	0.1539	0.1515	0.1492	0.1469	0.1446	0.1423	0.1401	0.1379	2	5	7	9	12	14	16	19	21
1.1	0.1357	0.1335	0.1314	0.1292	0.1271	0.1251	0.1230	0.1210	0.1190	0.1170	2	4	6	8	10	12	14	16	18
1.2	0.1151	0.1131	0.1112	0.1093	0.1075	0.1056	0.1038	0.1020	0.1003	0.0985	2	4	6	7	9	11	13	15	17
1.3	0.0968	0.0951	0.0934	0.0918	0.0901	0.0885	0.0869	0.0853	0.0838	0.0823	2	3	5	6	8	10	11	13	14
1.4	0.0808	0.0793	0.0778	0.0764	0.0749	0.0735	0.0721	0.0708	0.0694	0.0681	I	3	4	6	7	8	10	11	13
1.5	0.0668	0.0655	0.0643	0.0630	0.0618	0.0606	0.0594	0.0582	0.0571	0.0559	1	2	4	5	6	7	8	10	11
1.6	0.0548	0.0537	0.0526	0.0516	0.0505	0.0495	0.0485	00475	0.0465	0.0455	I	2	3	4	5	6	7	8	9
1.7	0.0446	0.0436	0.0427	0.0418	0.0409	0.0401	0.0392	0.0384	0.0375	0.0367	f	2	3	4	4	5	6	7	8
1.8	0.0359	0.0351	0.0344	0.0336	0.0329	0.0322	0.0314	0.0307	0.0301	0.0294	do	1	2	3	4	4	5	6	6
1.9	0.0287	0.0281	0.0274	0.0268	0.0262	0.0256	0.0250	0.0244	0.0239	0.0233	11	1/	2	2	3	4	4	5	5
2.0	0.0228	0.0222	0.0217	0.0212	0.0207	0.0202	0.0197	0.0192	8810.0	0.0183	0	ı	1	2	2	3	3	4	4
2.1	0.0179	0.0174	0.0170	0.0166	0.0162	0.0158	0.0154	0.0150	0.0146	0.0143	0	1	Dr.	2	2	2	3	3	4
2.2	0.0139	0.0136	0.0132	0.0129	0.0125	0.0122	0.0119	0.0116	0.0113	0.0110	0	41		1	2	2	2	3	3
2.3	0.0107	0.0104	0.0102							Pern	0	Y	l	1	1	2	2	2	2
				0.00990	0.00964	0.00939	0.00914			-	,426	Ġ.	8	10	13	15	18	20	23
								0.00889	0.00866	0.00842	2	5	7	9	12	14	16	16	21
2.4	0.00820	0.00798	0.00776	0.00755	0.00734						2	4	6	8	11	13	15	17	19
						0.00714	0.00695	0.00676	0.00657	0.00639	2	4	6	7	9	11	13	15	17
2.5	0.00621	0.00604	0.00587	0.00570	0.00554	0.00539	0.00523	0.00508	0.00494	0.00480	2	3	5	6	8	9	11	12	14
2.6	0.00466	0.00453	0.00440	0.00427	0.00415	0.00402	0.00391	0.00379	0.00368	0.00357	1	2	3	5	6	7	9	9	10
2.7	0.00347	0.00336	0.00326	0.00317	0.00307	0.00298	0.00289	0.00280	0.00272	0.00264	1	2	3	4	5	6	7	8	9
2.8	0.00256	0.00248	0.00240	0.00233	0.00226	0.00219	0.00212	0.00205	0.00199	0.00193	1	l	2	3	4	4	5	6	6
2.9	0.00187	0.00181	0.00175	0.00169	0.00164	0.00159	0.00154	0.00149	0.00144	0.00139	0	1	t	2	2	3	3	4	4
3.0	0.00135	0.00131	0.00126	0.00122	0.00118	0.00114	0,00111	0.00107	0.00104	0.00100	0	1	ı	2	2	2	3	3	4




Example / Contoh:

If $X \sim N(0, 1)$, then P(X > k) = Q(k)Jika $X \sim N(0, 1)$, maka P(X > k) = Q(k)

[Lihat halaman sebelah SULIT



Numbers in each row of the table are values on a t-distribution with (df) degrees of freedom for selected right-tail (greater-than) probabilities (p).



df/p	0.40	0.25	0.10	0.05	0.025	0.01	0.005	0.0005
1	0.324920	1.000000	3.077684	6.313752	12.70620	31.82052	63.65674	636.6192
2	0.288675	0.816497	1.885618	2.919986	4.30265	6.96456	9.92484	31.5991
3	0.276671	0.764892	1.637744	2.353363	3.18245	4.54070	5.84091	12,9240
4	0.270722	0.740697	1.533206	2.131847	2.77645	3.74695	4.60409	8.6103
5	0.267181	0.726687	1.475884	2.015048	2.57058	3.36493	4.03214	6.8688
6	0.264835	0.717558	1.439756	1.943180	2.44691	3.14267	3.70743	5.9588
7	0.263167	0.711142	1.414924	1.894579	2.36462	2.99795	3.49948	5.4079
8	0.261921	0.706387	1.396815	1.859548	2.30600	2.89646	3.35539	5.0413
9	0.260955	0.702722	1.383029	1.833113	2.26216	2.82144	3.24984	4.7809
10	0.260185	0.699812	1.372184	1.812461	2.22814	2.76377	3.16927	4.5869
11	0.259556	0.697445	1.363430	1.795885	2.20099	2.71808	3,10581	4.4370
12	0.259033	0.695483	1.358217	1.782288	2.17881	2.68100	3.05454	43178
13	0.258591	0.693829	1.350171	1.770933	2.16037	2.65031	3.01228	4.2208
14	0.258213	0.692417	1.345030	1,761310	2.14479	2.62449	2.97684	4.1405
15	0.257885	0.691197	1.340606	1.753050	2.13145	2.60248	2.94671	4.0728
16	0.257599	0.690132	1.336757	1.745884	2.11991	2.58349	2.92078	4.0150
17	0.257347	0.689195	1.333379	1.739607/	2.10982	2.56693	2.89823	3.9651
18	0.257123	0.688364	1.330391	1.734064	2.10092	2.55238	2.87844	3.9216
19	0.256923	0.687621	1.327728	1.729133	2.09302	2,53948	2.86093	3.8834
20	0.256743	0.686954	1.325341	1.724718	2.08596	2.52798//	2.84534	3.8495
21	0.256580	0.686352	1.323188	1.720743	2.07961	2.51765	2.83136	3.8193
22	0.256432	0.685805	1.321237	1.717144	2.07387	2.50832	2.81876	3.7921
23	0.256297	0.685306	1.319460	1.713872	2.06866	2.49987	2.80734	3.7676
24	0.256173	0.684850	1.317836	1.710882	2.06390	2.49216	2.79694	3.7454
25	0.256060	0.684430	1.316345	1.708141	2.05954	2.48511	2.78744	3.7251
26	0.255955	0.684043	1.314972	1.705618	2.05553	2.47863	2.77871	3.7066
27	0.255858	0.683685	1.313703	1.703288	2.05183	2.47266	2.77068	3.6896
28	0.255768	0.683353	1.312527	1.701131	2.04841	2.46714	2.76326	3.6739
29	0.255684	0.683044	1.311434	1.699127	2.04523	2.46202	2.75639	3.6594
30	0.255605	0.682756	1.310415	1.697261	2.04227	2.45726	2.75000	3.6460
Z	0.253347	0.674490	1.281552	1.644854	1.95996	2.32635	2.57583	3.2905
CI			80%	90%	95%	98%	99%	99.9%

